



## **Antimicrobial Resistance in Gram-Negative Rods Causing Bacteremia in Hematopoietic Stem Cell Transplant Recipients**

### **Intercontinental Prospective Study of the Infectious Diseases Working Party of the European Bone Marrow Transplantation Group**

Averbuch, Diana; Tridello, Gloria; Hoek, Jennifer; Mikulska, Malgorzata; Akan, Hamdi; Yanez San Segundo, Lucrecia; Pabst, Thomas; Özçelik, Tülay; Klyasova, Galina; Donnini, Irene; Wu, Depei; Gülbas, Zafer; Zuckerman, Tsila; Botelho de Sousa, Aida; Beguin, Yves; Xhaard, Aliénor; Bachy, Emmanuel; Ljungman, Per; de la Camara, Rafael; Rascon, Jelena; Ruiz Camps, Isabel; Vitek, Antonin; Patriarca, Francesca; Cudillo, Laura; Vrhovac, Radovan; Shaw, Peter J; Wolfs, Tom; O'Brien, Tracey; Avni, Batia; Silling, Gerda; Al Sabty, Firas; Graphakos, Stelios; Sankelo, Marja; Sengelojev, Henrik; Pillai, Srinivas; Matthes, Susanne; Melanthiou, Frederiki; Iacobelli, Simona; Styczynski, Jan; Engelhard, Dan; Cesaro, Simone

*Published in:*

Clinical Infectious Diseases

*DOI:*

[10.1093/cid/cix646](https://doi.org/10.1093/cid/cix646)

*Publication date:*

2017

*Document version*

Version created as part of publication process; publisher's layout; not normally made publicly available

*Document license:*

[CC BY-NC-ND](#)

*Citation for published version (APA):*

Averbuch, D., Tridello, G., Hoek, J., Mikulska, M., Akan, H., Yanez San Segundo, L., Pabst, T., Özçelik, T., Klyasova, G., Donnini, I., Wu, D., Gülbas, Z., Zuckerman, T., Botelho de Sousa, A., Beguin, Y., Xhaard, A., Bachy, E., Ljungman, P., de la Camara, R., ... Cesaro, S. (2017). Antimicrobial Resistance in Gram-Negative Rods Causing Bacteremia in Hematopoietic Stem Cell Transplant Recipients: Intercontinental Prospective Study of the Infectious Diseases Working Party of the European Bone Marrow Transplantation Group. *Clinical Infectious Diseases*, 65(11), 1819-1828. <https://doi.org/10.1093/cid/cix646>



UNIVERSITÀ  
DEGLI STUDI  
DI UDINE

## Università degli studi di Udine

### Antimicrobial resistance in Gram-negative rods causing bacteremia in hematopoietic stem cell transplant patients: intercontinental prospective

This is a pre print version of the following article:

#### *Original*

Antimicrobial resistance in Gram-negative rods causing bacteremia in hematopoietic stem cell transplant patients: intercontinental prospective study of Infectious Diseases Working Party of the European Bone Marrow Transplantation group / Averbuch, Diana; Tridello, Gloria; Hoek, Jennifer; Mikulska, Malgorzata; Akan, Hamdi; Yanez San Segundo, Lucrecia; Pabst, Thomas; Özçelik, Tülay; Klyasova, Galina; Donnini, Irene; Wu, Depei; Gülbaz, Zafer; Zuckerman, Tsila; Botelho De Sousa, Aida; Beguin, Yves; Xhaard, Aliénor; Bachy, Emmanuel; Ljungman, Per; De La Camara, Rafael; Rascon, Jelena; Ruiz Camps, Isabel; Vitek, Antonin; Patriarca, Francesca; Cudillo, Laura; Vrhovac, Radovan; Shaw, Peter; J. Wolfs, Tom; O'Brien, Tracey; Avni, Batia; Silling, Gerda; Al Sabty, Firas; Graphakos, Stelios; Sankelo, Marja; Sengeloev, Henrik; Pillai, Srinivas; Matthes, Susanne; Melanthiou, Frederiki; Iacobelli, Simona; Styczynski, Jan; Engelhard, Dan; Cesaro, Simone. - In: CLINICAL INFECTIOUS DISEASES. - ISSN 1058-4838. - STAMPA. - 65:11(2017), pp. 1819-1828.

*Publisher:*

#### *Published*

DOI:10.1093/cid/cix646

#### *Terms of use:*

The institutional repository of the University of Udine (<http://air.uniud.it>) is provided by ARIC services. The aim is to enable open access to all the world.

*Publisher copyright*

(Article begins on next page)



# Antimicrobial Resistance in Gram–Negative Rods Causing Bacteremia in Hematopoietic Stem Cell Transplant Recipients: Intercontinental Prospective Study of the Infectious Diseases Working Party of European Bone Marrow Transplantation Group

Diana Averbuch ✉ ✉, Gloria Tridello, Jennifer Hoek, Malgorzata Mikulska, Hamdi Akai, Lucrecia Yañez San Segundo, Thomas Pabst, Tülay Özçelik, Galina Klyasova, Irene Donnelly  
[Show more](#)

[Author Notes](#)

*Clinical Infectious Diseases*, cix646, <https://doi.org/10.1093/cid/cix646>

**Published:** 26 July 2017    **Article history** ▼

[PDF](#)   [Cite](#)   [Permissions](#)   [Share](#) ▼

## Abstract

### Background

This intercontinental study aimed to study gram–negative rod (GNR) resistance in hematopoietic stem cell transplantation (HSCT).

### Methods

GNR bacteremias occurring during 6 months post–HSCT (February 2014–May 2015) were prospectively collected, and analyzed for rates and risk factors for resistance to fluoroquinolones, noncarbapenem anti-*Pseudomonas*  $\beta$ -lactams (noncarbapenem  $\beta$ -lactams), carbapenems, and multidrug resistance.

### Results

resistance were higher in auto-HSCT patients in centers providing vs not providing fluoroquinolone prophylaxis ( $P < .01$ ). Resistance rates were higher in southeast and northwest Europe and similar in children and adults, excluding higher fluoroquinolone- and  $\beta$ -lactam/ $\beta$ -lactamase inhibitor resistance rates in allo-HSCT adults. Non-*Klebsiella* Enterobacteriaceae were rarely carbapenem resistant. Multivariable analysis revealed resistance risk factors in allo-HSCT patients: fluoroquinolone resistance: adult, prolonged neutropenia, breakthrough on fluoroquinolones; noncarbapenem resistance: hospital-acquired infection, breakthrough on noncarbapenems or other antibiotics (excluding fluoroquinolones, noncarbapenems, carbapenems), donor type; carbapenem resistance: breakthrough on carbapenem, longer hospitalization, intensive care unit, previous other antibiotic therapy; multidrug resistance: longer hospitalization, breakthrough on  $\beta$ -lactams,  $\beta$ -lactamase inhibitors, and carbapenems. Inappropriate empiric therapy and multidrug resistance were significantly more common in infections caused by resistant bacteria.

## Conclusions

Our data question the recommendation for fluoroquinolone prophylaxis and call for reassessment of local empiric antibiotic protocols. Knowledge of pathogen-specific resistance enables early appropriate empiric therapy. Monitoring of resistance is crucial.

---

**Keywords:** [antimicrobial resistance](#), [gram-negative rods](#), [hematopoietic stem cell transplantation](#), [bacteremia](#)

**Topic:** [antibiotics](#), [bacteremia](#), [carbapenem](#), [hematopoietic stem cell transplantation](#), [adult](#), [allopurinol](#), [child](#), [drug resistance](#), [microbial](#), [enterobacteriaceae](#), [fluoroquinolones](#), [lactams](#), [bacteria](#), [pathogenic organism](#), [gram-negative bacillus](#), [antibiotic resistance](#), [carbapenem resistance](#)

**Issue Section:** [Major Article](#)

studies, mainly limited to single centers or countries, describe rates and risk for specific resistant pathogens, such as extended-spectrum  $\beta$ -lactamase (ESBL)-producing Enterobacteriaceae or multidrug-resistant (MDR) bacteria in patients with cancer and HSCT recipients [10–13]. The majority, however, were performed in countries with high resistance rates in both the hospitalized and general populations [12–14]. The risk of infection is influenced by local factors, including prevalence of resistance, prophylactic practices, empiric treatment, and antimicrobial stewardship. Only a handful of studies focus on children and autologous HSCT (auto-HSCT) patients [12, 15, 16]. This study aimed to describe resistance rates and risk factors in GNR bacteremia in HSCT patients, based on intercontinental data.

## METHODS

---

### Study Design and Data Collection

This was a noninterventional prospective study. All patients in whom allogeneic HSCT (allo-HSCT) or auto-HSCT was performed during February 2014–March 2015 at the participating centers were included. Data on GNR bacteremia episodes that occurred from the initiation of the conditioning regimen until 6 months after HSCT were reported, including pathogen and its antimicrobial susceptibility, presence of risk factors, and mortality. This study was performed in accordance with the appropriate regulations in the participating countries including approval by the ethical committees as required, and registered at ClinicalTrials.gov (NCT02257931).

The primary endpoint was to determine the proportion of GNRs resistant to carbapenems and of noncarbapenem anti-*Pseudomonas*  $\beta$ -lactams (noncarbapenems), including

# Microbiological Workup

Guidelines used to determine isolates' susceptibility were by the European Committee for Antimicrobial Susceptibility Testing in 41 (64.1%) laboratories, Clinical and Laboratory Standards Institute in 19 (29.7%) laboratories, and 4 (6.2%) laboratories. Resistance to antibiotics was studied by in vitro sensitivity tests in the local laboratories using disk diffusion in 46 of 64 (71.9%) and/or minimum inhibitory concentration determination in 60 of 64 (93.8%) centers. Pathogens with intermediate susceptibility to antibiotics were considered resistant.

## Definitions

An MDR GNR was defined as bacteria resistant to  $\geq 1$  agent in  $\geq 3$  of the following categories: (1) broad-spectrum cephalosporins (ceftazidime or cefepime); (2) *Pseudomonas*  $\beta$ L/ $\beta$ LI; (3) carbapenems; (4) aminoglycosides; (5) fluoroquinolones. All *Stenotrophomonas maltophilia* strains were considered MDR. Extensively drug-resistant (XDR) bacteria defined if checked for susceptibility to all relevant antimicrobials and found nonsusceptible to  $\geq 1$  agent in all but  $\leq 2$  categories. Infections occurring  $>48$  hours since the hospitalization were considered hospital-acquired.

Neutropenia was defined as an absolute neutrophil count  $<500$  cells/ $\mu$ L.

Breakthrough bacteremia was defined as bacteremia developing during antimicrobial treatment (including fluoroquinolone prophylaxis) provided for  $\geq 48$  hours before obtaining the blood culture.

Geographic regions [4] included Northwest (Austria, Belgium, Denmark, Finland, France, Germany, Netherlands, Sweden, Switzerland, United Kingdom); South

The incidence of resistance was computed as a percentage, the denominator being the number of GNR pathogens and the numerator being the number of pathogens classified as “resistant.” Early mortality was computed as a percentage of deaths that occurred within 7 days after bacteremia on the episodes with a follow-up alive at the same time.

The relationship between resistance and the following risk factors was investigated:

- Background: sex, age at HSCT, underlying disease, myeloablative conditioning, fluoroquinolone prophylaxis.
- At the time of bacteremia: time since HSCT, duration of neutropenia, neutrophil recovery, duration of hospitalization, hospital-acquired infection, breakthrough bacteremia, graft-vs-host disease (GVHD), acute GVHD grade II–IV, veno-occlusive disease.
- Before bacteremia, within 1 month: urinary catheter,  $\geq 2$  weeks of steroids or other immunosuppressive treatment; within 3 months: any hospitalization, hospitalization in the intensive care unit (ICU), any bacteremia, previous antibiotic therapy, and number of antibiotic classes.

Differences between groups were tested using linear or logistic regression models using the generalized estimating equation method to take into account the dependence of observations nested by patient and center [18]. The same models were used to study all relationships between resistance and characteristics and prognostic factors. Variables showing significance from the univariate models entered a multivariable model. The results obtained from these analyses are considered as exploratory and hypothesis-generating.

A  $P$  value  $< .05$  was considered statistically significant. All  $P$  values are 2 sided. The analyses were performed using the statistical software SAS version 9.4.



Sixty-five HSCT centers from 25 countries reported data on 655 GNR episodes in 591 patients (1.1 episodes [range, 1–4] per patient) (Supplementary Table 1). Characteristics of the background patients and episodes are presented in [Table 1](#) and [2](#). Median time to develop GNR bacteremia was 8 (range, –15 to 183) days after HSCT. Fluoroquinolone prophylaxis was provided in 34 of 45 (75.6%) allo-HSCT adult centers, 22 of 49 (44.9%) auto-HSCT adult centers, and in 6 of 24 (25.0%) allo-HSCT pediatric centers, and 1 of 23 (8.7%) auto-HSCT pediatric centers.

**Table 1.**

Patient Characteristics (N = 591)

Characteristic	No. (%)
Age, y, median (range)	50.9 (0.3–77)
Children (<18 y)	79 (13.4)
Male sex	357 (60.4)
Underlying disease	
Acute leukemia	212 (35.9)
Lymphoma	148 (25.0)
Plasma cell disorders	109 (18.4)
Myelodysplastic/myeloproliferative syndromes	54 (9.1)
Chronic leukemia	8 (1.4)
Solid tumor	12 (2.0)

Unrelated	175 (48.9)
Matched related	114 (31.8)
Mismatched related	69 (19.3)
Stem cell source	
Peripheral blood	451 (76.4)
Bone marrow	101 (17.1)
Both bone marrow and peripheral blood	10 (1.7)
Umbilical cord blood	28 (4.7)
Conditioning (allogeneic HSCT)	
Myeloablative	241 (68.1)
Nonmyeloablative	113 (31.9)
Karnofsky score, median (range)	90 (0–100)
Disease status after HSCT	
Continued complete remission	317 (75.1)
Never in complete remission	79 (18.7)
Partial remission/stable disease	26 (6.2)
Graft-vs-host disease prophylaxis before bacteremia (allogeneic HSCT)	
Total provided	339

Data are presented as No. (%) unless otherwise indicated.

Abbreviations: HSCT, hematopoietic stem cell transplantation; MMF, mycophenolate mofetil.

View Large

Table 2.

Episode Characteristics

Characteristic	Transplant Type		Total (N = 655)
	Allogeneic (n = 414)	Autologous (n = 241)	
Neutropenia at the time of episode <500 cells/ μL	296 (71.5)	199 (82.9)	495 (75.7)
Neutropenia at the time of episode <100 cells/ μL	259 (73.4)	175 (77.8)	434 (75.1)
Duration of neutropenia <500 cells/μL, d, median (range)	7 (1–246)	2.5 (1–124)	5 (1– 246)
Duration of neutropenia <500 cells/μL <7 d, median (range)	153 (37.0)	189 (78.8)	342 (52.3)
Duration of neutropenia <100 cells/μL, d, median (range)	5 (1–80)	2 (1–28)	3 (1– 80)
Duration of neutropenia <100 cells/μL <7 d	176 (49.9)	171 (76.0)	347 (60.0)
Absolute neutrophil count recovered before bacteremia	147 (39.9)	24 (10.8)	171 (28.9)

Hospital-acquired bacteremia	355 (86.8)	215 (90.7)	570 (88.2)
Previous ICU hospitalization	28 (6.9)	4 (1.7)	32 (4.9)
Bacteremia developed during hospitalization in the ICU	17 (4.2)	0 (0.0)	17 (2.6)
Breakthrough bacteremia	303 (73.4)	98 (40.7)	401 (61.3)
Cephalosporins <sup>a</sup>	30 (7.3)	8 (3.3)	38 (5.8)
Anti- <i>Pseudomonas</i> βL/βLIs	65 (15.7)	16 (6.6)	81 (12.4)
Carbapenems <sup>b</sup>	54 (13.1)	7 (2.9)	61 (9.3)
Any β-lactam <sup>c</sup>	159 (38.5)	36 (14.9)	195 (29.8)
Fluoroquinolones	118 (28.6)	50 (20.7)	168 (25.7)
Aminoglycosides	22 (5.3)	5 (2.1)	27 (4.1)
Others <sup>d</sup>	79 (19.1)	13 (5.4)	92 (14.1)
Previous antibiotic therapy	309 (77.3)	104 (44.4)	413 (65.1)
Cephalosporins <sup>a</sup>	46 (11.5)	8 (3.4)	54

			(37.9)
Fluoroquinolones	121 (30.3)	45 (19.2)	166 (26.2)
Aminoglycosides	30 (7.5)	5 (2.1)	35 (5.5)
Others <sup>d</sup>	101 (25.3)	12 (5.1)	113 (17.8)
No. of antibiotic classes of previous antibiotic therapy <sup>e</sup>			
1 antibiotic class	86 (21.4)	36 (15.3)	122 (19.2)
2 classes	73 (18.2)	24 (10.2)	97 (15.2)
≥3 antibiotic categories	131 (32.6)	18 (7.7)	149 (23.4)
GVHD at the time of bacteremia (% of all allo-HSCT)	71 (17.3)		
Acute GVHD at the time of bacteremia (% of all allo-HSCT)	62 (15.1)		
Grade II–IV of acute GVHD before bacteremia (% of all allo-HSCT)	12 (4.5)		
Chronic GVHD (% of all allo-HSCT)	9 (2)		
Veno-occlusive disease at the time of bacteremia	12 (2.9)	3 (1.3)	15 (2.3)

Recent steroid treatment	142 (34.5)	53 (22.3)	195 (30.0)
Time since HSCT, d, median (range)	11 (–15 to 183)	7 (–8 to 177)	8 (–15 to 183)
Central line infection	138 (38.4)	73 (32.3)	211 (36.1)

Data are presented as No. (%) unless otherwise indicated.

Abbreviations: allo, allogeneic; auto, autologous;  $\beta$ L/ $\beta$ LI,  $\beta$ -lactam/ $\beta$ -lactamase inhibitor; GVHD, graft-versus-host disease; HSCT, hematopoietic stem cell transplant; ICU, intensive care unit.

<sup>a</sup> Cephalosporins: ceftazidime, cefepime, ceftriaxone, cefotaxime, cefuroxime, cefazolin.

<sup>b</sup> Carbapenems: meropenem, imipenem, ertapenem, doripenem.

<sup>c</sup> Any  $\beta$ -lactam: either cephalosporin, anti-*Pseudomonas*  $\beta$ L/ $\beta$ LI, carbapenem, or any other  $\beta$ -lactam antibiotic.


<sup>d</sup> Others: any antibiotics, excluding any  $\beta$ -lactam, fluoroquinolones, aminoglycosides, and prophylactic use of trimethoprim-sulfamethoxazole or dapsone.

<sup>e</sup> Previous antibiotic exposure to the following antibiotic classes was reported: (a) any cephalosporins; (b) anti-*Pseudomonas*  $\beta$ L/ $\beta$ LI; (c) carbapenems (meropenem, imipenem, ertapenem, doripenem); (d) fluoroquinolones (either treatment and prophylaxis); (e) aminoglycosides; (f) macrolides; (g) anti-*Staphylococcus aureus* glycopeptides (glycopeptides, linezolid, daptomycin); (h) antianaerobes (metronidazole, clindamycin); (j) tigecycline; (k) aztreonam; (l) other antibiotics.

[View Large](#)

## Microbiological Results

Seven hundred four GNRs included Enterobacteriaceae (514 [73%]),



Distribution of gram-negative pathogens. Other Enterobacteriaceae include *Enterobacter* spp (n = 10), *Klebsiella* spp (n = 24), *Citrobacter* spp (n = 12), *Serratia* spp (n = 6), *Proteus* spp (n = 3), *Raoultella* spp (n = 1). Other nonfermentative rods include *Acinetobacter* spp (n = 17), 1 each of *Sphingomonas paucimorosa*, *Shewanella putrefaciens*, *Sphingobacterium multivorum*, *Comamonas testosteroni*, *Ochrobactrum tritici*, *Ralstonia pickettii*, *Paracoccus yeei*, *Achromobacter xylosoxidans*. Other gram-negative rods include *Capnocytophaga* spp (n = 8), *Moraxella* spp (n = 4), *Aeromonas* spp (n = 3), *Campylobacter* spp (n = 1), *Rhizobium radiobacter* (n = 2), and 1 each of *Salmonella enteritidis*, *Haemophilus influenzae*, *Burkholderia cepacia*, *Elizabethkingia meningoseptica*.

## Resistance Rates

Half of GNRs were resistant to fluoroquinolones and to noncarbapenems, 18.2% were carbapenem resistant, and 35.2% were MDR ([Table 3](#)). One of 73 (1.4%) *Pseudomonas aeruginosa* and 4 of 14 (28.6%) *Acinetobacter* species were XDR. Resistance rates to other antibiotics are presented in [Table 3](#).

**Table 3.**

						Adu
Primary endpoints						
Fluoroquinolone	325/645 (50.4)	227/408 (55.6)*	98/237 (41.4)*	205/326 (62.9)**	22/82 (26.8)**	95/2 (41.7)
Noncarbapenem β-lactam	322/632 (50.9)	241/407 (59.2)*	81/225 (36.0)*	199/322 (61.8)	42/85 (49.4)	78/2 (36.4)
Carbapenem	127/688 (18.5)	105/443 (23.7)*	22/245 (8.9)*	80/349 (22.9)	25/94 (26.6)	20/2 (8.6)
Multidrug resistant	236/671 (35.2)	187/428 (43.7)*	49/243 (20.2)*	149/334 (44.6)	38/94 (40.4)	46/2 (19.9)
Secondary endpoints						
Anti- <i>Pseudomonas</i> cephalosporin	255/656 (38.9)	199/416 (47.8)	56/240 (23.3)	160/332 (48.2)	39/84 (46.4)	55/2 (24.0)
Anti- <i>Pseudomonas</i> βL/ βLI	230/627 (36.7)	177/399 (44.4)	53/228 (23.2)	146/310 (47.1)***	31/89 (34.8)***	51/2 (23.5)
Aminoglycoside	216/662 (32.6)	162/422 (38.4)	54/240 (22.5)	135/335 (40.3)	27/87 (31.0)	51/2 (22.4)
Colistin	22/358 (6.1)	18/250 (7.2)	4/108 (3.7)	12/205 (5.9)	6/45 (13.3)	4/10 (3.9)
Tigecycline	15/182 (8.2)	12/124 (9.7)	3/58 (5.2)	10/109 (9.2)	2/15 (13.3)	3/58 (5.2)

Data are presented as no./No. (%). The following differences were significant: the resistance rates of fluoroquinolones, noncarbapenems, carbapenems, and multidrug resistance were significantly higher



There was no strong correlation between the rates of GNR bacteremia and resistance to noncarbapenems and carbapenems per country; and no correlation between fluoroquinolone resistance and bacteremia rate per center (Supplementary 1A–C).

## Resistance Rates According to Pathogens

Fluoroquinolone resistance was significantly more frequent among Enterobacteriaceae (57.2% vs 30.7%;  $P < .0001$ ); carbapenem resistance (50.8.4%;  $P < .0001$ ), and multidrug resistance (46.6% vs 31.9%;  $P = .001$ ) in nonfermentative rods ([Table 4](#)).

**Table 4.**

Resistance Rates According to Pathogens

Pathogens	Resistance to			
	Fluoro-quinolones	Noncarbapenem β-Lactams	Carbapenem	Multidrug Resistant
<i>Escherichia coli</i>	185/283 (65.4)	140/281 (49.8)	7/301 (2.3)	81/290 (27.9)
<i>Klebsiella pneumoniae</i>	71/111 (63.9)	79/118 (66.9)	31/124 (25.0)	63/121 (52.1)
<i>Enterobacter</i> spp	8/39 (20.5)	19/39 (48.7)	3/41 (7.3)	9/39 (23.1)
Other Enterobacteriaceae	7/41 (17.1)	9/43 (20.9)	2/45 (4.4)	5/44 (11.4)

<i>Stenotrophomonas maltophilia</i>	6/22 (27.3)	14/16 (87.5)	34/34 (100)	34/34 (100)
Other nonfermentative rods	4/22 (18.2)	13/20 (65.0)	5/21 (23.8)	7/22 (31.8)
Total nonfermentative rods	46/150 (30.7) <sup>a</sup>	68/138 (49.3)	82/161 (50.9) <sup>a</sup>	76/163 (46.6) <sup>b</sup>

Data are presented as no./No. (%).

<sup>a</sup> Significant differences between Enterobacteriaceae and nonfermentative rods ( $P < .0001$ ).

<sup>b</sup> Significant differences between Enterobacteriaceae and nonfermentative rods ( $P = .001$ ).

[View Large](#)

Six of 31 (19%) *S. maltophilia* isolates were resistant to trimethoprim-sulfamethoxazole.

One-third of noncarbapenem-resistant, and half of MDR pathogens, were carbapenem resistant (Supplementary Table 2). Six of 33 (18.2%) carbapenem-resistant *P. aeruginosa* isolates were susceptible to both cephalosporins and (Supplementary Table 3); 1 of 29 (3.5%) was colistin resistant.

Among carbapenem-resistant Enterobacteriaceae, 15 of 41 (36.6%) were resistant to amikacin, 16 of 41 (39.0%) to gentamicin, 4 of 41 (9.8%) to both aminoglycosides, 32 of 40 (78%) to fluoroquinolones, 5 of 17 (29.4%) to tigecycline, and 6 of 33 (18.2%) to colistin.

## Resistance Rate According to Hematopoietic Stem Cell Transplantation

HSCT patients (Supplementary Tables 4 and 5).

### Resistance Rate in Children Versus Adults

Similar resistance rates were observed in children and adults. Only the resistance rate to fluoroquinolones and  $\beta$ L/ $\beta$ LI was significantly higher in adults compared with children following allo-HSCT ( $P < .0001$  and  $P = .048$ , respectively) (Table 5); the difference was significant in southeast countries only.

### Geographical Distribution of Resistance Rates

There was a wide distribution in the resistance rates between countries (Supplementary Table 6). Resistance rates were more frequent in the southeast than in the northwest European region: for noncarbapenems (266/481 [55.3%] vs 32/118 [27.6%],  $P < .0001$ ); fluoroquinolones (270/496 [54.4%] vs 44/121 [36.4%];  $P = .002$ ); carbapenems (109/526 [20.7%] vs 6/123 [4.9%];  $P < .0001$ ); and multidrug resistance (201/514 [39.1%] vs 16/118 [13.6%];  $P < .0001$ ).

### Risk Factors for Resistance

Univariate analysis of risk factors for resistant GNRs in allo-HSCT and autologous HSCT recipients is presented in Supplementary Tables 4 and 5. Multivariable analysis was not possible in allo-HSCT patients only (Table 5), as rate of resistance was low in autologous HSCT patients.

**Table 5.**

Multivariate Analysis of Risk Factors for Resistance in Gram-Negative Rods in Allogeneic Transplantation

Risk Factor	Fluoroquinolone-Resistant GNRs	Noncarbapenem $\beta$ -Lactam-Resistant GNRs	Carbapenem-Resistant GNRs	Multidrug-Resistant GNRs
-------------	--------------------------------	--	---------------------------	--------------------------

Breakthrough bacteremia on fluoroquinolones	7.26 (3.84–13.72)	<.0001		NS		NS	
Age (adults)	3.87 (2.07–7.25)	<.0001		NS		NS	
Hospital-acquired infection		NS	2.09 (1.08–4.03)	.029		NS	
Breakthrough bacteremia on βL/βLI		NS	2.45 (1.14–5.23)	.02		NS	2.3 (1.1–4.4)
Breakthrough bacteremia on cephalosporins		NS	6.36 (1.68–24.07)	.01		NS	
Previous antibiotic therapy with other antibiotics <sup>a</sup>		NS	2.59 (1.30–5.15)	.007	2.07 (1.15–3.73)	.016	
Longer duration of current hospitalization before episode		NS		NS	1.01 (1.00–1.02)	.041	1.0 (1.0–1.0)
Current bacteremia developed during hospitalization in the ICU		NS		NS	3.92 (1.18–3.03)	.026	
Breakthrough		NS		NS	9.08	<.0001	3.4

unrelated donor

(2.22  
4.35)

Abbreviation:  $\beta$ L/ $\beta$ LI, anti-*Pseudomonas*  $\beta$ -lactam  $\beta$ -lactamase inhibitor; CI, confidence interval; C negative rod; ICU, intensive care unit; NS, not significant; OR, odds ratio.

<sup>a</sup> Other antibiotics: any antibiotic therapy excluding  $\beta$ -lactams, fluoroquinolones, aminoglycosi

[View Large](#)

To study the association between fluoroquinolone prophylaxis and resistance rates in GNRs compared resistance rates in GNRs cultured during the period when prophylaxis was provided (in neutropenic patients without empiric or targeted antibiotics therapy) in centers that do and do not provide prophylaxis.

The rate of fluoroquinolone-resistant GNRs was higher in centers providing prophylaxis (79% vs 50%,  $P = .001$  in allo-HSCT; 74% vs 25%,  $P < .001$  in auto-HSCT). The rate of noncarbapenem-resistant (36% vs 13%;  $P = .002$ ) and MDR (46% vs 8%;  $P < .001$ ) bacteria was higher in auto-HSCT patients in centers providing prophylaxis (Supplementary Table 7).

## Outcome

The 7-day mortality was 38 of 589 (6.5%). The mortality according to resistance pattern was 9% vs 2% ( $P = .002$ ) in episodes caused by noncarbapenem resistant vs sensitive GNRs; 18% vs 4% ( $P < .001$ ) in those carbapenem resistant vs sensitive and 11% vs 4% ( $P = .002$ ) in MDR vs non-MDR. Inappropriate empiric therapy was provided for 124 of 586 (21.2%) GNRs; being 101 of 270 (37.4%) for noncarbapenem resistant vs 12 of 274 (4.4%) noncarbapenem susceptible bacteria; 62 of 97 (63.9%) for carbapenem resistant vs 61 of 481 (12.7%) carbapenem susceptible; and 46 (46.2%) for MDR vs 30 of 368 (8.1%) non-MDR GNRs ( $P < .001$  for all).

resistance.

The emerging resistance challenges antibacterial prophylaxis policy and complicates empiric and targeted treatment choices. Benefit of fluoroquinolone prophylaxis was demonstrated in a country with a baseline resistance of approximately 20% in GNRs from the community and medical departments. In our study, 39% of GNRs causing community-acquired infections in allo- and 63% in auto-HSCT were fluoroquinolone resistant (Supplementary Table 5). High rates of fluoroquinolone resistance may be the price to pay for low rates of bacteremia in centers providing prophylaxis. We could not, however, demonstrate a correlation between the rates of bacteremia and fluoroquinolone resistance (Supplementary Figure 1A). Doubts regarding the benefits of fluoroquinolone prophylaxis are accompanied by concern about increased rates of resistance to other antibiotics following exposure to fluoroquinolones. No association between fluoroquinolone prophylaxis and the increase in MDR bacteria was shown in a meta-analysis of studies published up to 2005 [21]. Later published studies, however, correlated exposure to fluoroquinolones with increase in infections resulting from fluoroquinolone-resistant, ESBL-producing, carbapenem-resistant, and MDR pathogens [14, 22, 23]. In our study, in auto-HSCT patients, treatment in centers providing fluoroquinolone prophylaxis, breakthrough on fluoroquinolones, and previous exposure to fluoroquinolones (Supplementary Tables 5 and 7) were associated with resistance to noncarbapenems and MDR. The risk could, however, be influenced by other factors, as outpatient vs inpatient transplantation settings, which were not reported in our study. The benefits and potential risks of fluoroquinolone prophylaxis must be carefully assessed, especially in centers with high fluoroquinolone resistance rates among GNRs [24].

Empiric treatment with noncarbapenem  $\beta$ -lactams, or even carbapenems [25], may be inappropriate in centers with high resistance rates to these antibiotics and may lead to increased mortality [2, 13]. In our study, approximately 40% of community-

for empiric therapy are tricky because of significant geographical variations. Practical decisions on empiric therapy must be based on continuously updated data concerning local resistance patterns and bacteremia rates [26]. We are thus unable to recommend the specific resistance rate threshold that indicates change in empiric therapy protocol as our study was not designed to answer this question. While the proportion of resistant bacteria was high in some centers, its impact on patient outcome can be low if GNR bacteremia is rare (Supplementary Figure 1 and 1C).

Resistance to multiple antibiotics complicates the targeted therapy choice. Susceptibility to tigecycline, polymyxins, fosfomycin, and aminoglycosides was promptly reported in the centers with carbapenem-resistant infections, as *K. pneumoniae* bacteria were resistant to these last-resort antibiotics in our and other studies [27]. Eighteen percent of carbapenem-resistant *P. aeruginosa* was susceptible to cephalosporins and  $\beta$ L/ $\beta$ LI in our study and, in contrast to carbapenem-resistant Enterobacteriaceae, can be treated with these agents, especially administered with high-dose prolonged infusion [28].

Knowledge of pathogen-specific resistance patterns can help direct appropriate empiric therapy following rapid bacterial identification by matrix-assisted laser desorption/ionization–time of flight (MALDI-TOF), prior to susceptibility results. Carbapenem monotherapy should be appropriate on identification of non-*K. pneumoniae* Enterobacteriaceae, which are usually carbapenem susceptible. Carbapenem/colistin with or without aminoglycoside combination should, however, be considered for *K. pneumoniae* or *Acinetobacter* pending susceptibility results, as a significant proportion of them are carbapenem resistant. Streamlining of antibiotic therapy, of course, should be performed when susceptibilities are available.

Data on resistance rates in post-HSCT children are scarce [12, 16, 29]. Surprisingly, resistance rates were mostly similar in children and in adults in our study.



Information about resistance rates and risk factors in auto-HSCT patients is limited and mainly concerns fluoroquinolones [15, 30], as bacteremia is relatively less frequent in auto-HSCT patients [31]. Certain factors predisposing to resistant infections—such as breakthrough bacteremia and prolonged neutropenia—are more frequent in auto-HSCT vs allo-HSCT patients (Table 2). Although total resistance rates were higher in allo-HSCT patients, resistance rates in community-acquired infections were similar. These data reinforce the importance of monitoring antibiotic susceptibilities in auto-HSCT patients.

Several studies reported risk factors for cephalosporin-resistant, ESBL-producing, or MDR bacteria in cancer and HSCT patients [10, 12, 13, 16, 22, 29, 32]. Only a handful describe risk factors for carbapenem-resistant infections in populations involving but not limited to transplant patients, mainly for *K. pneumoniae*. In addition, from other studies, we found that breakthrough on noncarbapenems does not predispose to carbapenem resistance in allo-HSCT patients [8, 14]. Noncarbapenem  $\beta$ -lactams and carbapenems are both recommended for empiric treatment in neutropenic patients [25], half of whom have neither microbiological nor clinical evidence of infection. Limitation of carbapenem treatment, de-escalation to a narrower spectrum regimen following culture results, and shortening treatment duration are recommended by European Conference on Infections in Leukaemia guidelines to slow development of carbapenem resistance without increasing mortality [1].

Geographical differences in resistance rates are striking. In certain countries, we found higher resistance rates than those previously reported in the literature. In France, for example, 34.4% of GNRs were noncarbapenem resistant, compared with 14% third-generation cephalosporin resistance among Enterobacteriaceae carriage in hematological malignancies and post-HSCT patients during 2003–2010, which may be explained by increased resistance rates over time [5]. Information about resistance rates in each country may, of course, be skewed by local epidemiology.



significantly higher resistance rates in southeast as compared with northwest Europe. This correlates with lower rates of antimicrobial resistance among general population [36], which is probably explained by lower consumption of systemic antibacterials in the community, as well as in veterinary medicine in northwest Europe [37, 38].

The multidrug resistance rate (~30%) among Enterobacteriaceae and *P. aeruginosa* in our study was within that reported in cancer and HSCT patients [11–13, 39]. The rate of cotrimoxazole resistance among *S. maltophilia*, the drug of choice for this MDR pathogen, was higher in our study (19%) compared to 4%–10% in other studies [5, 29]. Susceptibility of *Acinetobacter* species in cancer and transplant patients has been infrequently studied [5, 29, 39]; XDR *Acinetobacter* infections have not previously been reported. In our study, a third of *Acinetobacter* species were XDR.

We demonstrated higher mortality rates in infections caused by resistant bacteria, similar to other studies [8, 12]. Analysis of mortality risk factors is beyond the scope of this manuscript.

Our study has limitations. GNR resistance rates and patterns in our study are likely influenced by the epidemiology of countries with more participating centers. Antimicrobial susceptibility data were incomplete for some GNRs. Resistance to certain agents, such as tigecycline, could be overestimated, as susceptibility was likely checked in bacteria resistant to other treatment options, or reflecting local epidemiology. We could, however, demonstrate resistance rates to salvage treatments among their main targets, harder-to-treat pathogens. We did not identify important risk factors for resistance: history of recent residency in another country (information unavailable) and prior colonization with resistant GNRs (not all participating centers performed colonization screening).

The study has important strengths. It is the first intercontinental prospective

studies, we analyze risk factors for different resistance patterns and pathogens rather than focusing on specific bacteria or mechanisms.

In conclusion, the problem of antibiotic resistance is worrying in all HSCT patients, including subgroups of children and auto-HSCT recipients. It is associated with inappropriate empiric therapy and increased mortality. Benefits of fluoroquinolone prophylaxis and the approach to empiric therapy should be reassessed and adapted to continuous monitoring of the local bacteremia rates and susceptibility data of the infecting pathogens. Knowledge of pathogen-specific resistances enables the choice of appropriate empiric therapy.

## Supplementary Data

---

Supplementary materials are available at *Clinical Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted supplementary materials are not copyedited and are the sole responsibility of the authors, all questions or comments should be addressed to the corresponding author.

## Notes

---

**Acknowledgments.** This is a multicenter study. We are deeply indebted to the members of the Infectious Diseases Working Party of the European Bone Marrow Transplantation group for fruitful discussions and to all collaborators: Czech Republic: Alexandra Jungova, Petra Keslova, Milan Navratil; France: Aziza Azzou Nedhif; Emmanuelle Nicolas; Germany: Arnold Ganser; Greece: Anastasia Pappa; Israel: Elizabeth Neparstek; Italy: Claudio Annaloro, Elisabetta Calore, Katia Codeluppi, Edoardo Lanino, Francesco Lanza, Katia Perruccio, Roberto Rainone.

Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

## References

---

1. Passweg JR , Baldomero H , Bader P et al. Use of haploidentical stem cell transplant continues to increase: the 2015 European Society for Blood and Marrow Transplantation survey report. *Bone Marrow Transplant* 2017; 52:811–7.  
[Google Scholar](#)   [CrossRef](#)   [PubMed](#)
2. Girmenia C , Viscoli C , Piciocchi A et al. Management of carbapenem resistant *Klebsiella pneumoniae* infections in stem cell transplant recipients: an Italian multidisciplinary consensus statement. *Haematologica* 2015; 100:e373–6.  
[Google Scholar](#)   [CrossRef](#)   [PubMed](#)
3. Averbuch D , Cordonnier C , Livermore DM et al. Targeted therapy against multi-resistant bacteria in leukemic and hematopoietic stem cell transplant recipients: guidelines from the European Conference on Infections in Leukemia (ECIL-4, 2011). *Haematologica* 2013; 98:1836–47.  
[Google Scholar](#)   [CrossRef](#)   [PubMed](#)
4. Mikulska M , Viscoli C , Orasch C et al. Fourth European Conference on Infections in Intensive Care and Hematology (ECIL-4), a joint venture of EBMT, EORTC, ICHS, ELN and ESGICH/ESCMID . Antimicrobial use and resistance in bacteraemias among adult and paediatric haematology and cancer patients. *J Infect* 2014; 68:321–31.  
[Google Scholar](#)   [CrossRef](#)   [PubMed](#)
5. Bousquet A , Malfuson JV , Sanmartin N et al. An 8-year survey of strains identified in blood cultures in a clinical haematology unit. *Clin Microbiol Infect* 2014; 20:O7–12.

7. Girmenia C , Rossolini GM , Picciocchi A et al. Gruppo Italiano Trapianto Midollo Osseo GITMO . Infections by carbapenem-resistant *Klebsiella pneumoniae* in SCT recipients: a nationwide retrospective survey in Italy. *Bone Marrow Transplant* 2015; 50:282–8.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

8. Satlin MJ , Cohen N , Ma KC et al. Bacteremia due to carbapenem-resistant Enterobacteriaceae in neutropenic patients with hematologic malignancies. *J Infect Dis* 2007; 195:7336–45.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

9. Mikulska M , Del Bono V , Raiola AM et al. Blood stream infections in allogeneic hematopoietic stem cell transplant recipients: reemergence of gram-negative rods and increasing resistance. *Biol Blood Marrow Transplant* 2009; 15:47–53.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

10. Vehreschild MJ , Hamprecht A , Peterson L et al. A multicentre cohort study on colonization and infection with ESBL-producing Enterobacteriaceae in high-risk patients with haematological malignancies. *J Antimicrob Chemother* 2014; 69:3387–92.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

11. Trecarichi EM , Pagano L , Candoni A et al. HeMABIS Registry—SEIFEM Group, Italy: epidemiology and antimicrobial resistance data for bacterial bloodstream infections in adult patients with hematologic malignancies: an Italian multicentre prospective survey. *Microbiol Infect* 2015; 21:337–43.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

12. Caselli D , Cesaro S , Ziino O et al. Infection Study Group of the Associazione Italiana Ematologia Oncologia Pediatrica (AIEOP) . Multidrug resistant *Pseudomonas aeruginosa* infection in children undergoing chemotherapy and hematopoietic stem cell transplantation. *Haematologica* 2010; 95:1612–5.

14. Trecarichi EM , Pagano L , Martino B et al. Haematologic Malignancies Associated Bloodstream Infections Surveillance (HEMABIS) registry—Sorveglianza Epidemiologica Infezioni Fungine in Emopatie Maligne (SEIFEM) group, Italy . Bloodstream infection by *Klebsiella pneumoniae* in onco-hematological patients: clinical impact of carbapenem resistance in a multicentre prospective survey. *Am J Hematol* 2016; 91:1076–81.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

15. Wang CH , Chang FY , Chao TY et al. Characteristics comparisons of bacteremia in allogeneic and autologous hematopoietic stem cell-transplant recipients with levofloxacin prophylaxis and influence on resistant bacteria emergence. *J Microbiol Immunol Infect* 2016. doi:10.1016/j.jmii.2016.02.003.

16. Satwani P , Freedman JL , Chaudhury S et al. A multicenter study of bacterial bloodstream infections in pediatric allogeneic hematopoietic cell transplantation recipients: the impact of acute gastrointestinal graft-versus-host disease. *Biol Blood Marrow Transplant* 2016; 23:642–7.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

17. Magiorakos AP , Srinivasan A , Carey RB et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012; 18:268–81.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

18. Lipsitz SR , Fitzmaurice GM , Orav EJ , Laird NM . Performance of generalized estimating equations in practical situations. *Biometrics* 1994; 50:270–8.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

19. Bucaneve G , Micozzi A , Menichetti F et al. Gruppo Italiano Malattie Ematologiche dell'Adulto (GIMEMA) Infection Program . Levofloxacin to prevent bacterial infections in patients with cancer and neutropenia. *N Engl J Med* 2005; 353:977–87.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

22. Liss BJ , Vehreschild JJ , Cornely OA et al. Intestinal colonisation and blood stream infections due to vancomycin-resistant enterococci (VRE) and extended-spectrum lactamase-producing Enterobacteriaceae (ESBLE) in patients with haematological oncological malignancies. *Infection* 2012; 40:613–9.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

23. Liu CY , Lai YC , Huang LJ et al. Impact of bloodstream infections on outcome and influence of prophylactic oral antibiotic regimens in allogeneic hematopoietic SC recipients. *Bone Marrow Transplant* 2011; 46:1231–9.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

24. Kim JH , Goulston C , Zangari M , Tricot G , Boyer MW , Hanson KE . Impact of a chemotherapy-related antibacterial prophylaxis on bacteremia and hospitalization rates following outpatient autologous peripheral blood stem cell transplantation for multiple myeloma. *Trans Infect Dis* 2014; 16:421–9.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

25. Freifeld AG , Bow EJ , Sepkowitz KA et al. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2011; 52:e56–93.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

26. Averbuch D , Orasch C , Cordonnier C et al. European guidelines for empirical antimicrobial therapy for febrile neutropenic patients in the era of growing resistance: summary of the 2011 4th European Conference on Infections in Leukemia. *Haematologica* 2013; 98:1826–35.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

27. Pagano L , Caira M , Trecarichi EM et al. Carbapenemase-producing *Klebsiella pneumoniae*

29. Averbuch D , Avaky C , Harit M et al. Non-fermentative gram-negative rods bacteria in children with cancer: a 14-year single-center experience. *Infection* 2017; 45:327–33.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

30. Satlin MJ , Vardhana S , Soave R et al. Impact of prophylactic levofloxacin on rate of bloodstream infection and fever in neutropenic patients with multiple myeloma undergoing autologous hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant* 2015; 21:1808–14.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

31. Bock AM , Cao Q , Ferrieri P , Young JA , Weisdorf DJ . Bacteremia in blood or marrow transplantation patients: clinical risk factors for infection and emerging antibiotic resistance. *Biol Blood Marrow Transplant* 2013; 19:102–8.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

32. Patriarca F , Cigana C , Massimo D et al. Risk factors and outcomes of infections by multidrug-resistant gram-negative bacteria in patients undergoing hematopoietic transplantation. *Biol Blood Marrow Transplant* 2017; 23:333–9.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

33. Mokart D , Slehofer G , Lambert J et al. De-escalation of antimicrobial treatment in neutropenic patients with severe sepsis: results from an observational study. *Intensive Care Med* 2014; 40:41–9.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)

34. Blennow O , Ljungman P , Sparrelid E , Mattsson J , Remberger M . Incidence, risk factors and outcome of bloodstream infections during the pre-engraftment phase in 521 allogeneic hematopoietic stem cell transplantations. *Transpl Infect Dis* 2014; 16:106–14.

[Google Scholar](#)      [CrossRef](#)      [PubMed](#)



ropa.eu/en/eaad/Documents/antibiotics-EARS-Net-summary-2016.pdf. Accessed 2017.

37. European Centre for Disease Prevention and Control. Geographical distribution of antimicrobial consumption. Available at: <http://ecdc.europa.eu/en/healthtopics/antibiotic-resistance-and-consumption/antimicrobial-consumption/esac-net-data-base/Pages/geo-distribution-consumption.aspx>. Accessed 7 April 2017.
38. European Medicines Agency. Sales of veterinary antimicrobial agents in 26 EU/EEA countries in 2013. Available at: [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Report/2015/10/WC500195687.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Report/2015/10/WC500195687.pdf). Accessed 7 April 2017.
39. Kim SB , Min YH , Cheong JW et al. Incidence and risk factors for carbapenem- and multidrug-resistant *Acinetobacter baumannii* bacteremia in hematopoietic stem cell transplantation recipients. *Scand J Infect Dis* 2014; 46:81–8.

[Google Scholar](#)

[CrossRef](#)

[PubMed](#)

© The Author 2017. Published by Oxford University Press for the Infectious Disease Society of America. All rights reserved. For permissions, e-mail: [journals.permissions@oup.com](mailto:journals.permissions@oup.com).

## Supplementary data

---

[supplementary\\_materials](#)

– docx file

## Comments

---



## Email alerts

[New issue alert](#)

[Advance article alerts](#)

[Article activity alert](#)

---

[Receive exclusive offers and updates  
from Oxford Academic](#)

## More on this topic

Trends in Antimicrobial Resistance in Health  
Care–Associated Pathogens and Effect on  
Treatment

Response to Rituximab-Based Therapy and Risk  
Factor Analysis in Epstein Barr Virus–Related  
Lymphoproliferative Disorder After  
Hematopoietic Stem Cell Transplant in Children  
and Adults: A Study From the Infectious  
Diseases Working Party of the European Group  
for Blood and Marrow Transplantation

An Antimicrobial Stewardship Program's Impact

*Escherichia coli* Sequence Type ST131 as the  
Major Cause of Serious Multidrug-Resistant *E.*  
*coli* Infections in the United States

## Related articles in

Antibiotics Versus Surgical Therapy for Uncomplicated Appendicitis: Contrasting Interpretations of Data.

Reply to: The letter to the editor "Antibiotics vs. Surgical Therapy for Uncomplicated Appendicitis: Contrasting Interpretations of Data" by McAnena et al.

Relationship between Dental Procedures and Shunt Infections in Hydrocephalic Patients: A Narrative Review.

## Citing articles via

Google Scholar

CrossRef

**Latest** | **Most Read** | **Most Cited**

Outbreak of invasive infections from subtype *emm26.3* group A *Streptococcus* among homeless adults—Anchorage, Alaska, 2016–2017

Breast implant Q fever as a source of in hospital transmission

Isolation and identification of *R. raoultii* in human cases: a surveillance study in three medical centers in China

*Salmonella enterica* serotype 4,[5],12:i:- in swine in the United States Midwest: an emerging multidrug resistant clone

[Author Guidelines](#)

[Advertising and Corporate Services](#)

[Facebook](#)

[Journals Career Network](#)

[Twitter](#)

# Clinical Infectious Diseases

Online ISSN 1537-6591

Print ISSN 1058-4838

Copyright © 2017 Infectious Diseases Society of America

[About Us](#)

[Contact Us](#)

[Careers](#)

[Help](#)

[Access & Purchase](#)

[Rights & Permissions](#)

[Open Access](#)

## Connect

[Join Our Mailing List](#)

[OUPblog](#)

[Twitter](#)

[Facebook](#)

[YouTube](#)

[Tumblr](#)

## Resources

[Authors](#)

[Librarians](#)

[Societies](#)

[Sponsors & Advertisers](#)

[Press & Media](#)

[Agents](#)

